

Kills at least 99.9999% of spores in 2 minutes

in dirty conditions<sup>1</sup>

**CLINELL SPORICIDAL WIPES** 

# THE MOST CLINICALLY PROVEN DISINFECTANT WIPE IN THE WORLD

## Effective against dry biofilms

Clinell Sporicidal Wipes have shown effective removal of dry biofilm pathogens as well as preventing transfer and regrowth<sup>2</sup>.

## Safe and easy to use

Premeasured dose is activated by wetting with water - no more dilution errors. No toxic fumes are produced upon activation so they are safe to use next to patients.

## The most clinically proven wipe in the world

Clinell Sporicidal Wipes kill most known pathogenic microorganisms with a non-selective action. With a proven 6 log spore kill, in two minutes under dirty conditions, it is proven to outperform other leading sporicidal wipes<sup>1,3</sup>.

# Faster & more effective in highly soiled conditions

More effective than two-step cleaning and disinfection with chlorine<sup>4</sup>. Allows both cleaning and disinfection, remaining effective in highly soiled conditions unlike chlorine based products<sup>5,6</sup>.

## Compatible with most surfaces

With nearly neutral pH, Clinell Sporicidal Wipes can be used on most non-invasive medical devices and equipment without corrosion.

## Patented formula

Clinell Sporicidal Wipes are CE certified class IIa medical devices that contain a unique powder combination to deliver peracetic acid on demand with just the addition of water.

# WHY CHOOSE CLINELL SPORICIDAL WIPES?

Improved cleaning and disinfection of room surfaces decreases the risk of healthcare associated infections<sup>7</sup>.

## Effective sporicidal agents

Enhanced environmental cleaning with sporicidal agents of rooms housing *Clostridioides difficile* infected patients is warranted<sup>8</sup>. Key measures to prevent *C. difficile* transmission include correct cleaning and disinfection of hospital room surfaces daily and at discharge. For effective disinfection of *C. difficile*, a sporicidal product plus correct practices are essential<sup>9,10</sup>.

### SPORICIDAL AND ANTI-BIOFILM

Quarternary Ammonium Compounds (QACs) are termed sporistatic<sup>3</sup> meaning they only inhibit spore germination and/or outgrowth. Clinell Sporicidal Wipes generate the oxidising agent peracetic acid, a well-known sporicidal agent. Clinell Sporicidal Wipes also demonstrate efficacy against dry biofilms<sup>2</sup>.

#### **TOXICOLOGICAL HAZARDS**

Chlorine solutions, made from tablets and powders, are hazardous to both the user and the patients: emitting toxic fumes and by-products. Chlorine has been shown to cause respiratory, nasal, skin and eye complaints<sup>12</sup>.

### **CHLORINE IS INEFFECTIVE**

One-step disinfectant wipes are more effective than two-step cleaning and disinfection with chlorine<sup>4</sup>. When preparing chlorine solutions, concentrations may be incorrect resulting in a disinfectant that is weak and ineffective and toxic to staff, patients and materials.



Biofilm formation provides protection and nutrients for microorganisms, allowing them to survive in hostile environments<sup>13</sup>.



Chlorine vapour causes lung damage, eye irritation, nose and throat damage, coughs and shortness of breath<sup>12</sup>.



Clinell Sporicidal Wipes are more effective at reducing the presence of Multi-Drug Resistant Organisms than two-step cleaning and disinfection with chlorine<sup>4</sup>.

# Chronic inhalation of chlorine can increase the risk of lung fibrosis<sup>12</sup>

Peracetic acid generating wipes activated by water, for surface disinfection and cleaning of non-invasive medical devices.



## **Clinell Sporicidal Wipes**

Containing patented technology, Clinell Sporicidal Wipes are designed for use on all surfaces of non-invasive medical devices. Upon activation with water, they generate peracetic acid levels proven to kill most microorganisms including dry biofilm pathogens.

Clinell Sporicidal Wipes are a high level disinfectant wipe highly effective against *C. difficile* spores. They clean and disinfect, providing a more effective and safe alternative to two-step detergent cleaning followed by chlorine solution disinfection<sup>3-6</sup>.

#### **ANTIMICROBIAL ACTIVITY**

Powerful disinfecting composition using a pH optimised peracetic acid and hydrogen peroxide combination, generated from sodium percarbonate and tetraacetylethylenediamine.

High power oxidative kill against all microorganisms including nonenveloped viruses, bacterial endospores and dry biofilms<sup>2</sup>.

Conforms to EN1276, EN13704, EN14348, EN14476, EN14561, EN14562, EN14563 and EN16615.

\* Independently tested in Professor Maillard's laboratory at Cardiff University. Test reports available on request.

BACTERIA	TEST	CONTACT TIME
Acinetobacter baumannii	EN13727	10 sec
Escherichia coli	EN13727	10 sec
Pseudomonas aeruginosa	EN13727	10 sec
	EN14561	1 min
Staphylococcus aureus	EN13727	10 sec
	EN14561	1 min
Enterococcus hirae	EN13727	10 sec
	EN14561	1 min
Klebsiella pneumoniae (ESBL)	EN13727	10 sec
Enterococcus faecalis	EN13727	10 sec
Dry Biofilms	ASTM E2967-15	Wiping 10 sec, contact 2 min
SPORES		
Clostridium difficile	EN13704	2 min
	ASTM E2362-09	2 min
Bacillus subtilis	EN13704	2 min
MYCOBACTERIA		
Mycobacterium terrae	EN14348	2 min
Mycobacterium avium	EN14348	2 min
FUNGI		
Candida auris	EN13624	10 sec
VIRUSES		
Adenovirus	EN14476	1 min
Human coronavirus (MERS-CoV)	EN14476	1 min
Canine parvovirus	EN14675	2 min



Dry before activation, each wipe consists of 2 non-woven fabrics bonded together with a unique powder combination in the middle layer.



Water activates the powder to generate high levels of hydrogen peroxide and peracetic acid.

PRODUCT INFORMATION	UNIT OF ISSUE	ORDER CODE	NHSSC
Sporicidal Wipes*	Pack of 25	CS25	VJT113
Sporicidal Wall Mounted Dispenser	for CS25	CSD	-



This unique method of application ensures consistent efficacy, with no dilution errors.

# Class Ila Medical Device

USE DISINFECTANT SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE. ALWAYS FOLLOW THE MEDICAL DEVICE MANUFACTURER'S CLEANING AND DISINFECTION GUIDELINES PRIOR TO USE OF THE WIPE.

# PROBLEMS WITH EXISTING SOLUTIONS

Irregular spray patterns lead to areas of low biocide concentration. Wet wipes always deliver the correct dose.

## Sprays and wipes

When using sprays and dry wipes, it is common to end up with variable concentrations of biocides<sup>15</sup>. Impregnated wet wipes are significantly better at removing microbial bioburden than when using a spray and dry wipe<sup>16</sup>.

Low concentrations of biocides can cause microorganisms to develop cross-resistance to antibiotics<sup>17,18</sup>.

Provided there is proper use of efficacious surface disinfectants, avoiding low concentrations of biocides, present scientific data does not suggest that resistance problems will emerge<sup>19,20</sup>. Wet wipes have the advantage of delivering a premeasured dose of biocides, avoiding the risk of resistance.

# Sprays provide uneven coverage.



Low biocide concentration may select for resistance.





## Two chamber sprays

Due to instability of the active biocide, or to prevent degradation over time, dual chamber sprays function by mixing two solutions within the spray head. It is common to use sprays at an angle, to shoot down onto the surface, meaning more solution will enter the head from one chamber than the other. This creates an uneven mix which can lead to suboptimal concentration of biocides.

In a laboratory setting, biocides may not be tested from dual chamber spray bottles – hence test data may not reflect practical, in-use conditions. In addition to affecting the efficacy of the biocide, inconsistent mixing increases wastage – reducing number of uses and resulting in increased cost.

The final mixed solution can build up over time and clog the nozzle. Blocked nozzles lead to even greater inconsistency of coverage than usual, resulting in low concentrations of biocides, which can allow microorganisms to develop cross-resistance<sup>17</sup>.



Wet wipes are significantly better at removing microbial bioburden than spray and dry wipes<sup>4</sup>

The molecular structure of the dry wipe can alter the formulation of the disinfectant by binding certain components and rendering them inactive<sup>21</sup>.

## Do dry wipes affect your disinfectant?



Dry wipes made of either synthetic or natural materials can bind the biocide, changing the concentration of active disinfectant released<sup>21</sup>.

This means that the liquid coming off the dry wipe may not be the same concentration as the liquid that went into it. For example, the combination of quaternary ammonium compounds (QACs) with an inappropriate type of fabric will effectively abolish its antimicrobial activity<sup>22</sup>.



Inactive biocide comes off the dry wipe.



Active biocide comes off the wet wipe.

Efficacy testing of disinfectant sprays and solutions has been performed on the liquid.

Efficacy testing of wet wipes is performed on the solution released by the wipe, and the wet wipe itself, so you know the end product exactly conforms to efficacy testing data.



Ineffective against microorganisms



Dry wipes can interfere with the action of hospital disinfectants<sup>21</sup>

# Pre-impregnated wet wipes are proven to increase staff compliance<sup>23</sup>

# CHLORINE SOLUTIONS

If a used wipe is plunged back into a bucket of chlorine, organic matter can be introduced. This accelerates breakdown of the active disinfectant.

## Double dipping

When a used wipe is dipped back into the bucket, organic matter is introduced, increasing breakdown of the active disinfectant.

The incompatibility of cloths and chlorine-based products has been clearly demonstrated<sup>11</sup>. Chlorine solution is tested in the laboratory for its effectiveness, however testing is not done on solution in conjunction with the cloth, but only on the solution by itself.

Shown to be more effective than twostep detergent cleaning followed by chlorine disinfection, Clinell Sporicidal Wipes contain detergent to clean as they disinfect whilst remaining effective in dirty conditions<sup>1,3</sup>.



Over diluted solutions are weak and ineffective.







Under diluted solutions are toxic and can cause damage to materials.



Organic matter binds to chlorine, reducing efficacy.

# DRY BIOFILMS

Dry biofilms pose significant challenges in clinical settings because they protect organisms contained within them, they are difficult to eradicate and are invisible to the naked eye<sup>13</sup>.

## Biofilms within healthcare settings

#### MICROORGANISM ACCUMULATION

Microorganisms accumulate and attach to surfaces forming communities known as biofilms<sup>24</sup>. It is thought that biofilm formation provides protection and nutrients for microorganisms allowing them to survive in hostile environments<sup>13</sup>.

#### DIFFICULT TO REMOVE

Biofilms are difficult to remove and survive decontamination methods previously thought to be effective in environmental decontamination<sup>25</sup>. The presence of dry biofilms harbouring bacterial pathogens is virtually universal on commonly used items in healthcare<sup>2</sup>. Many existing disinfectants do not demonstrate efficacy against dry biofilms.

# FOUND ON COMMONLY USED HEALTHCARE ITEMS:

- Patient folders
- Clipboards
- Chairs
- Commodes
- Sanitising bottles
- Trolleys
- Keyboard keys

## Measuring efficacy against dry biofilms

In collaboration with Cardiff University we have developed a method that has enabled measurement of the efficacy of Clinell Sporicidal Wipes against dry biofilms based on transferability, regrowth and removal<sup>26</sup>.

#### Transferability

Determines how well pathogens from dry biofilm can transfer to other surfaces post-wiping.

### Regrowth

Measures the time needed for a treated biofilm to regrow.

### Removal

Log reduction gives the number of bacteria that are removed following wiping.



## Clinell's industry leading aftercare and training services

Including a revolutionary training kit, specialised nurse trainers, educational materials, customised posters and dispenser installation.

Contact us on: 020 7993 0030 or info@clinell.com.

#### REFERENCES

- 1. Hospital Infection Research Laboratory. Sporicidal Efficacy Test. 2007. http://clinell.com/wp-content/ uploads/pdf/Clinell%20Sporicidal%20-%20 Efficacy%20Test.pdf.
- Ledwoch et al. Beware Biofilm! Dry biofilms containing bacterial pathogens on multiple healthcare surfaces; a multicentre study. J Hosp Infect 2018. In Press.
- Siani H et al. Efficacy of "sporicidal" wipes against Clostridium difficile. Am J Infect Control. 2011;39(3):212-8.
- Siani H et al. Impact of antimicrobial wipes compared with hypochlorite solution on environmental surface contamination in a health care setting: A doublecrossover study. Am J Infect Cont 2018. In Press.
- Humphreys PH et al. A systematic evaluation of a peracetic-acid-based high performance disinfectant. J Infect Prev 2013;14(4):126-31.
- Doan et al. Clinical and cost effectiveness of eight disinfection methods for terminal disinfection of hospital isolation rooms contaminated with Clostridium difficile 027. J Hosp Infect. 212; 82(2):114-21.
- 7. Weber DJ et al. The role of the surface environment in healthcare-associated infections. Curr Op Infect Dis. 2013 1;26(4):338-44.
- Weber et al. Role of the environment in the transmission of Clostridium difficile in health care facilities. Am J Infect Contr. 2013; 31;41(5):S105-10.
- Cadnum JL et al. Transfer of Clostridium difficile Spores by non-sporicidal wipes and improperly used hypochlorite wipes practice + product = perfection. Infect Cont Hosp Epidemiol. 2013; 34;(04):441-2.

- Maillard J-Y. Innate resistance to sporicides and potential failure to decontaminate. J Hosp Infect. 2011 31;77(3):204-9.
- Goldsmith M et al. Adsorption of available chlorine and quaternary by cotton and wool fabrics from disinfecting solutions. Appl Microbiol. 1954;2(6):360.
- Hoyle GW, Svendsen ER. Persistent effects of chlorine inhalation on respiratory health. Ann N Y Acad Sci. 2016; 1378(1):33-40.
- Otter JA et al. Surface-attached cells, biofilms and biocide susceptibility: implications for hospital cleaning and disinfection. J Hosp Infect. 2015;89(1):16-27.
- 14. Carter Y, Barry D. Tackling C difficile with environmental cleaning. Nurs Times. 2011; 107(36): 22-5.
- 15.Sattar SA, Maillard J-Y. The crucial role of wiping in decontamination of high-touch environmental surfaces: review of current status and directions for the future. Am J Infect Control. 2013; 31;41(5)S97-104
- Panousi MN et al. Evaluation of alcohol wipes used during aseptic manufacturing. Letters in Applied Microbiology. 1954;2(6):230.
- Maillard JY, Denyer SP. Emerging bacterial resistance following biocide exposure: should we be concerned? Chimica Oggi. 2009; 27(3): 26-8.
- 18.Karatzas KA et al. Phenotypic and proteomic characterization of multiply antibiotic-resistant variants of Salmonella enterica serovar Typhymurium selected following exposure to disinfectants. Appl Environ Microbiol. 2008 1;74(5):1508-16.
- 19.Gebel J et al. The role of surface disinfection in infection prevention. GMS Hyg Infect Cont. 2013;8(1)

- 20.Meyer B, Cookson B. Does microbial resistance of adaptation to biocides create a hazard in infection prevention and control? J Hosp Infect. 2010 30;76(3): 200-5.
- 21.Bloss R et al. Adsorption of active ingredients of surface disinfectants depends on the type of fabric used for surface treatment. J Hosp Infect. 2010; 75(1):56-61.
- 22.Engelbrecht K et al. Decreased activity of commercially available disinfectants containing quaternary ammonium compounds when exposed to cotton towels. Am J Infect Control. 2013;41(10):908-11.
- 23.Wiemken TL et al. The value of ready-to-use disinfectant wipes: compliance, employee time, and costs. Am J Infect Control. 2014; 31;42(3):329-30.
- 24.Donlan RM, Costerton JW. Biofilms: Survival Mechanisms of Clinically Relevant Microorganisms. Clin Microbiol Rev. 2002;15(2):167-93.
- 25.Vickery K et al. Presence of biofilm containing viable multiresistant organisms despite terminal cleaning on clinical surfaces in an intensive care unit. J Hosp Infect. 2012 80(1): 52-55.
- 26.Cardiff University. Transfer and Regrowth Test for the Evaluation of Anti-biofilm activity of Clinell Sporicidal Wipe. 2018. Full report can be found at clinell.com.



innovative solutions professional support

**GAMA Healthcare Ltd.**, 2 Regal Way, Watford, Hertfordshire, WD24 4YJ, UK. T: +44 (0)20 7993 0030 E: info@gamahealthcare.com www.gamahealthcare.com

